

AD-~~XXXXXXXXXX~~ A262 389
|||||

(2)

GL-TR 90-0310

Rational Approximation and Remote Temperature Sensing

DTIC
ELECTE
MAR 16 1993
S C D

Steven J. Leon

Southeastern Massachusetts University
Mathematics Department
North Dartmouth, MA 02747

20000920272

20 November 1990


**Reproduced From
Best Available Copy**

Final Report
Period Covered: October 1, 1989 - September 30, 1990

Approved for public release; distribution unlimited


GEOPHYSICS LABORATORY
AIR FORCE SYSTEMS COMMAND
UNITED STATES AIR FORCE
HANSCOM AIR FORCE BASE, MASSACHUSETTS 01731-5000


93-05376



98 3 15 044

This technical report has been reviewed and is approved for publication.


JEAN I.F. KING
Contract Manager
Atmospheric Sciences Division


ROBERT A. McCLATCHEY, Director
Atmospheric Sciences Division

This report has been reviewed by the ESD Public Affairs Office (PA) and is releasable to the National Technical Information Service (NTIS).

Qualified requestors may obtain additional copies from the Defense Technical Information Center. All others should apply to the National Technical Information Service.

If your address has changed, or if you wish to be removed from the mailing list, or if the addressee is no longer employed by your organization, please notify PL/TSI, Hanscom AFB, MA 01731-5000. This will assist us in maintaining a current mailing list.

Do not return copies of this report unless contractual obligations or notices on a specific document requires that it be returned.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0189

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 20 November 1990	3. REPORT TYPE AND DATES COVERED Final Report 1 Oct. 1989-30 Sept. 1990
----------------------------------	------------------------------------	---

4. TITLE AND SUBTITLE Rational Approximation and Remote Temperature Sensing	5. FUNDING NUMBERS PE 61102F PR 2310 TA G8 WU AI
--	--

6. AUTHOR(S) Steven J. Leon	Contract F19628-90-K-0008
--------------------------------	---------------------------

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Southeastern Massachusetts University Mathematics Department North Dartmouth, MA 02747	8. PERFORMING ORGANIZATION REPORT NUMBER
--	---

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Geophysics Laboratory Hanscom AFB, MA. 01731-5000 Contract Manager: Jean I. F. King/LY	10. SPONSORING / MONITORING AGENCY REPORT NUMBER GL-TR-90-0310
---	--

11. SUPPLEMENTARY NOTES

12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited	12b. DISTRIBUTION CODE
---	------------------------

13. ABSTRACT (Maximum 200 words) In recent years there have been a number of papers dealing with methods for remote temperature sensing based on applying transform theory to the radiative transfer equation. In particular, methods based on optical measure theory method have the advantage that they do not require computations of numerical derivatives. In this paper we present techniques for approximating the radiance profile based on rational interpolation and nonlinear least squares data fitting. Optical measure theory permits the inversion of non-Laplacian exponential kernels. The models developed are easily extended to include generalized exponential weighting functions. We also introduce the concept of total nonlinear least squares data fitting.
--

14. SUBJECT TERMS Remote temperature sensing, optical measure theory, differential inversion, radiative transfer, nonlinear least squares, rational approximation.	15. NUMBER OF PAGES 12
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	16. PRICE CODE
18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED
20. LIMITATION OF ABSTRACT SAR	

NSN 7540-01-280-5500

Standard Form 298 Rev. 2-89
Prescribed by ANSI Std. Z39-18
298-102

RATIONAL APPROXIMATION AND REMOTE TEMPERATURE SENSING

Steven J. Leon

Department of Mathematics
Southeastern Massachusetts University
North Dartmouth, Massachusetts 02747

Accession For	
NTIS	CRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced <input type="checkbox"/>	
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

1 INTRODUCTION

In recent years there have been a number of papers dealing with methods for remote temperature sensing based on applying transform theory to the radiative transfer equation. King, Hohlfeld, and Kilian (1989) have shown that using differential inversion it is possible to successfully determine temperature profiles from measurements of the upwelling intensities in the atmosphere. In this research project an alternative transform method based on optical measure theory is developed. Previous work on this method has been carried out by by King and Leon (1989,1990) and by Leon (1990). The optical measure theory method has the advantage that it does not require any computations of numerical derivatives. Instead the radiance profile is approximated by a rational function. The Planck intensity is then determined as the inverse transform of the rational function.

In Section 2 we present techniques for approximating the radiance profile based on rational interpolation and nonlinear least squares data fitting. Optical measure theory permits the inversion of non-Laplacian exponential kernels. Consequently, in Section 3, we extend the mathematical models to include generalized exponential weighting functions. In Section 4 we discuss alternative nonlinear data fitting techniques. In particular we introduce the

concept of total nonlinear least squares data fitting. Some test results are presented in Section 5 and conclusions follow in Section 6.

2 RATIONAL APPROXIMATION OF RADIANCE PROFILES

Using radiative transfer theory one can relate the Planck intensity to the upwelling intensity of the atmosphere. This relationship can be expressed in terms of an integral equation of the first kind. Specifically the relationship is given by

$$R(\hat{p}) = \int_0^{\infty} B(p) \mathcal{W}(p/\hat{p}) dp/p \quad (1)$$

where $\mathcal{W}(p/\hat{p})$ is a kernel weight function that peaks at $p = \hat{p}$ and $R(\hat{p})$ denotes the radiance of the wavelength channel whose weight function peaks at $p = \hat{p}$.

If we set

$$\mathcal{W}(z) = ze^{-z} \quad \text{and} \quad s = \frac{1}{\hat{p}}$$

then equation (1) can be expressed as a Laplace transform and consequently we can solve for the Planck function

$$B(p) = \mathcal{L}^{-1} \left(\frac{R(\frac{1}{s})}{s} \right)$$

We can think of $B(p)$ and $R(\hat{p})$ as transform pairs. Once a representation for R has been decided upon, then B can be determined analytically as an inverse Laplace transform.

The H -function inversion theory of Chandrasekhar, (1950) suggests that R should be represented as a rational function. If we set

$$R(\hat{p}) = d_1 + d_2\hat{p} + \cdots + d_{j+1}\hat{p}^j + \sum_{i=1}^l \frac{w_i}{1 + c_i\hat{p}} \quad (2)$$

then

$$\frac{R(\frac{1}{s})}{s} = \frac{d_1}{s} + \frac{d_2}{s^2} + \cdots + \frac{d_{j+1}}{s^{j+1}} + \sum_{i=1}^l \frac{w_i}{s + c_i}$$

and

$$B(p) = d_1 + d_2p + \cdots + \frac{d_{j+1}}{j!} p^j + \sum_{i=1}^l w_i e^{-c_i p} \quad (3)$$

The coefficients in equation (2) can be determined from the data by rational interpolation or by nonlinear least squares. Denote the radiance value corresponding to \hat{p}_i by R_i for $i = 1, \dots, m$. If one represents the radiance function as a rational function of the form

$$R(\hat{p}) = \frac{a_1 \hat{p}^n + a_2 \hat{p}^{n-1} + \dots + a_{n+1}}{\hat{p}^l - a_{n+2} \hat{p}^{l-1} - \dots - a_m} \quad (n + l = m - 1)$$

then R will interpolate (\hat{p}_i, R_i) if and only if

$$a_1 \hat{p}_i^n + a_2 \hat{p}_i^{n-1} + \dots + a_{n+1} + a_{n+2} \hat{p}_i^{l-1} R_i + \dots + a_m R_i = \hat{p}_i^l R_i \quad (4)$$

$i = 1, \dots, m$

The coefficients a_i are determined by solving the linear system (4). The residue form of R can be computed as an inverse convolution and from this one can determine the coefficients for equations (2) and (3). Generally these equations need only have a small number of components. It is possible to get a good fit with $j = 1, 2$ or 3 and $l = 1$ or 2 . If one takes $j = 2$ and $l = 2$, it is possible that for some data sets the interpolating function will have a positive pole, ($c_i < 0$), even though this is impossible for an actual radiance profile. When this happens the weight w_i corresponding to the pole will be much smaller than the weight corresponding to the other rational component. Thus if one sets $w_i = 0$, then the resulting function will give a good approximation to the data points.

In practice one can obtain a better approximation to the radiance data and avoid the problem of positive poles by using nonlinear least squares. The rational interpolating function can be taken as starting approximation for an iteratively computed nonlinear least squares fit to the data of the form

$$R(\hat{p}) = x_1 + x_2 \hat{p} + \dots + x_{j+1} \hat{p}^j + \sum_{i=1}^l \frac{x_{j+1+i}}{1 + x_{j+1+l+i} \hat{p}} \quad (5)$$

Initially the x_i coefficients are determined from the coefficients of the rational interpolating function of the form (2). Whenever the interpolating function has a positive pole, i.e., a coefficient $c_i < 0$, one sets the corresponding coefficient $x_{j+1+l+i}$ in (5) equal to 1 and the weight x_{j+1+i} can be set equal to 0.




Figure 1. Nonlinear least squares radiance function for set of TOVS data

The choice of a rational function to represent the radiance data is motivated by the physics of the atmosphere. In practice the nonlinear least squares approximation gives a very good fit to the radiance data. The coefficients of the rational function can be used to determine a Planck function of the form (3).

In Figure 1, a least squares fit of the form (5) is given for a typical data set obtained from the NOAA TIROS Operational Vertical Sounder (TOVS).

3 GENERALIZED EXPONENTIAL INVERSION

We can obtain a natural generalization of our previous results by taking a more general form for the weight function. If we set

$$W(z) = W_k(z) = \gamma_k z \exp_k(z)$$

where

$$\gamma_k = \frac{1}{k^{1/k} \Gamma\left(\frac{k+1}{k}\right)}$$

and

$$\exp_k(z) = \exp(-z^k/k)$$

then it follows that

$$R(\hat{p}) = \int_0^\infty B(p) W_k(p/\hat{p}) \frac{dp}{p}$$

If $R(\hat{p})$ is of the form (2), then the Planck intensity $B(p) = B_k(p)$ will be of the form

$$B_k(p) = d_1 + d_2 p + \cdots + \frac{d_{j+1}}{j!} p^j + \sum_{i=1}^l w_i \exp_k(c_i p) \quad (6)$$

Note that equations (3) and (6) are the same in the case $k = 1$.

4 TOTAL NONLINEAR LEAST SQUARES

A number of algorithms have been developed for nonlinear least squares fitting. Two algorithms using gradient methods to minimize the residual functions were developed specifically for rational fits of the form (5). These algorithms require good starting approximations and may converge to a local rather than a global minimum. The standard Nelder-Meade Simplex Algorithm, however, does seem to give satisfactory results even though convergence is rather slow. It may be possible to speed up convergence by taking some sort combination of the Nelder-Meade and gradient methods.

It should be noted that the standard nonlinear least squares techniques all seek to minimize the residual errors for the dependent variable. For the type of radiance fitting problem considered here, the values of the independent variable \hat{p} are also determined from satellite data and may involve some error. In fact, early experiments seem to indicate that the computed Planck function is more sensitive to perturbations in the \hat{p}_i values than to changes in the R_i values. In order to reflect this in the data fitting, algorithms have been devised for total nonlinear least squares fits. In this process the residual is defined in terms of the sum of the squares of the distances of the points (\hat{p}_i, R_i) to tangent lines to the curve (5) at the points $(\hat{p}_i, R(\hat{p}_i))$. Indeed a parameter t has been incorporated into the data fitting algorithms to specify

which type of fit is used. If $t = 0$, then the usual nonlinear least squares fit is used to minimize the residuals of the dependent variable. If $t = 1$, then a total least squares fit is computed. If $t = 2$, then a rational least squares fit is computed to minimize the residuals of the independent variable. For values of t between 0 and 2, intermediate types of fits are computed.

5 TEST RESULTS

Equation (1) is an integral equation of the first kind and consequently the problem of finding a solution is ill-posed. In addition to testing the optical measure theory method we also tested some of the standard methods used for solving integral integral equations of the first kind.

The standard techniques involve discretizing the equation and adding regularization conditions. The problem is then translated into a linear system of equations with quadratic constraints. The constrained problem can be solved using the methods proposed by Gander, (1981). However, when these techniques were applied to data sets obtained from the NOAA TIROS Operational Vertical Sounder (TOVS), the resulting matrices had very low numerical rank. Consequently, even with regularization conditions we were unable to obtain meaningful results. On the other hand, the transform methods have smoothness constraints and additional structure built in. The rational form (2) can be computed in a numerically stable manner and the coefficients define a unique Planck function.

The interpolation and nonlinear least squares algorithms have been tested extensively on simulated data and on the TOVS data. The algorithms are numerically stable. Figures 2 through 5 represent Planck functions that were generated using the same data set as in Figure 1. Figure 2 shows a semilog plot of the Planck function. The pressures are given in millibars on the vertical axis. Figure 3 shows the Planck function for the data set derived by generalized exponential inversion with $k = 0.8$. Figure 4 shows the Planck function derived by generalized exponential inversion with $k = 1.2$. Figure 5 shows all three Planck functions plotted as functions of p on the same axis system.




Figure 2. Planck intensity derived from set of TOVS data




Figure 3. Generalized exponential inversion with $k = 0.8$




Figure 4. Generalized exponential inversion with $k = 1.2$




Figure 5. $k = 0.8, 1, 1.2$ inverses

The question that still is outstanding is how to choose j , the degree of the polynomial component of the fit, and how to choose l , the number of hyperbolic components. Although it is reasonable that both of these numbers should be relatively small, it is not possible to give a definitive answer as to the optimal values of j and l based solely on the TOVS data sets and simulated data. It is anticipated that these questions will be answered when future data sets with more sensing channels in the appropriate ranges become available.

6 CONCLUSIONS

Although transfer theory can be used to relate the Planck intensity to the upwelling intensity in the atmosphere, the relation is expressed in the form of an integral equation of the first kind. Such equations are ill-posed and consequently do not have unique numerical solutions. Adding regularization conditions does not solve the problem. For data sets like those obtained from the TOVS data, there is not enough information to determine the Planck function using matrix methods with appropriate smoothness constraints. More assumptions relative to the physics of the atmosphere must be added. In this regard the transform methods based on optical measure theory seem to work well. The key step is the representation of the radiance profile by a rational function. This can be accomplished in a numerically stable manner using nonlinear least squares fitting algorithms.

References

- [1] S. Chandrasekhar. *Radiative Transfer*. Oxford University Press, London, 1950.
- [2] Walter Gander. Least squares with a quadratic constraint. *Numerische Mathematik*, 36, 1981.
- [3] Robert G. Hohlfeld, James C. Kilian, Thomas W. Drueding, and John F. Ebersole. *Novel Methodology for Application of Adaptive Systems Techniques to DMSP Remote Temperature Sensing*. Technical Report 88-0323, AFGL, Hanscom AFB, Massachusetts 01731, 1988. ADA209975
- [4] Robert G. Hohlfeld, Ronald F. Wachtmann, and James C. Kilian. *Application and Evaluation of a Differential Inversion Technique for Re-*

remote Temperature Sensing. Technical Report 88-0138, AFGL, Hanscom AFB, Massachusetts 01731, 1988. ADA199155

- [5] J. I. F. King, R. G. Hohlfeld, and J. C. Kilian. Application and evaluation of a differential inversion technique for remote temperature sensing. *Met Atmos. Phys.*, 41, 1989.
- [6] Jean I. F. King. Theory and application of differential inversion to remote sensing. In A. Deepak et al., editors, *Advances in Remote Sensing*, A. Deepak Publishing, Hampton, Virginia, 1985.
- [7] Jean I. F. King and Steven J. Leon. A smart algorithm for nonlinear interpolation and noise discrimination. In A. Deepak et al., editors, *Advances in Remote Sensing Retrieval Methods*, A. Deepak Publishing, Hampton, Virginia, 1989.
- [8] Jean I. F. King and Steven J. Leon. Smooth profile inferencing from noisy radiance data using optical measure theory. In *Fifth Conference on Satellite Meteorology and Oceanography, Preprint Volume*, American Meteorological Society, Boston, MA, 1990.
- [9] Steven J. Leon. *Algorithms for Interpolating and Approximating Radiance Profiles*. AFGL Technical Report, AFGL, Hanscom AFB, Massachusetts 01731, 1990. GL-TR-90-0071
- [10] Steven J. Leon. Matrix and other methods for solving integral equations of the first kind with applications to remote temperature sensing. *J. of Linear Algebra and its Applications*, vol. 150. May 1991.

**END
FILMED**

DATE:

4-93

DTIC